

APESIN spray F

Alcohol-based rapid disinfection, liquid – perfume-free

Hard non-porous small surface disinfection | Ready-to-use



Technical Information



Product profile

- APESIN spray F is a fast-acting disinfectant against bacteria, yeasts, enveloped viruses (limited virucidal) and viruses (limited virucidal PLUS).
- Effectiveness tested under high organic load (disinfectant cleaner).
- Surfaces that come into direct contact with food do not need to be rinsed after using the product. This saves one work step.
- The product dries without leaving residues.
- APESIN Spray F is free of chlorine, aldehydes and phenols.
- VAH and IHO listed.

Area of application

- APESIN spray F can be applied to aluminum surfaces.
- Do not use on alcohol-sensitive materials (e.g. PLEXIGLAS®).
- **PT02** - Indoor:
 - Health care facilities and pharmaceutical and cosmetic industry, for example patient-near surrounding, working areas/desks, general equipment (with the exception of surfaces that come into contact with food).
- **PT04** - Health care facilities and in food industry, for example food preparation and handling in kitchens/restaurants.
- Authorisation number EU-0027671-000

Application note

- Surfaces should always be visibly clean before disinfection.
- Ready-to-use surface disinfectant at room temperature (20±2 °C).
- The entire surface to be disinfected is wetted by pouring or spraying from a short distance and subsequently thoroughly wiped with a cloth. The amount of product should be sufficient (max. 50 ml/m²) to keep the surface wet during the contact time.
- **PT02** - Application rate: Exposure time: - for the control of bacteria, yeasts and enveloped viruses: 60 sec - for the control of viruses (limited spectrum virucidal activity): 5 min.
 - Number and timing of application: A reasonable frequency of disinfection in a patient's room is 1-2 per day. Maximum number of applications is 6 per day. No safety intervals need to be considered between the application phases.
- **PT04** - Application rate: Minimum exposure time: for the control of bacteria and yeasts at 20°C: 60 sec.
 - Number and timing of application: the products can be used as often as necessary. A reasonable frequency in kitchens is 1-2 per day. No safety intervals need to be considered between the application phases.

Sales units:


- 716672 - 10 x 750 ml
- 1114867 - 10 x 750 ml
- 1314929 - 750 ml
- 716671 - 10 x 750 ml
- 1116058 - 2x 5 l



Application and Dosing




Follow below instructions.



concentrated


Disinfection

Suitable for areas with specific hygiene requirements.



Surface disinfection:

Wet the surface completely.




Toilet disinfection



concentrated

Ideal for application on surfaces in kitchen areas.



APESIN spray F

PT 04 Target organism (s)	Dosage / Dilution	minimum exposure time
Bacteria	Ready-to-use	60 s
Yeasts		

PT 02 Target organism (s)	Dosage / Dilution	minimum exposure time
Bacteria	Ready-to-use	60 s
Yeasts		
Viruses (Virucidal activity against enveloped viruses)	Ready-to-use	5 min
Viruses (Limited spectrum virucidal activity)		

The results have been substantiated to the competent authority by testing the following standards:
 EN 1276, EN 1650, EN 13697, EN 13727, EN 13624, EN 13697 mod., EN 16615, EN 14476 (+A1, 2016).
 Viruses (Limited spectrum virucidal activity) spectrum of efficacy: Enveloped viruses and Norovirus, Rotavirus and Adenovirus

VAH - Certificate		minimum exposure time
Surface disinfection in medical areas, bacterial and yeasticidal activity	with mechanical action dirty conditions	5 Min.

Ingredient declaration

Ingredients per 100g: 40g ISOPROPYL ALCOHOL, 25g PROPYL ALCOHOL

Handling, storage and environmental instructions

Product strictly reserved for a professional use. For all information regarding handling, storage conditions and disposal, please see the Safety Data Sheet. Use biocidal products with care. Always read label and product information before use.

Company certificates of Management Systems

