


APESIN spray F



Alcohol-based rapid disinfection, liquid – perfume-free

- Hard non-porous small surface disinfection
- Ready-to-use

Product profile

- APESIN spray F is a fast-acting disinfectant against bacteria, yeasts, enveloped viruses (limited virucidal) and viruses (limited virucidal PLUS).
- Surfaces that come into direct contact with food do not need to be rinsed after using the product. This saves one work step.
- The product dries without leaving residues.
- APESIN Spray F is free of chlorine, aldehydes and phenols.
- VAH and IHO listed.

Area of application

- APESIN spray F can be applied to aluminum surfaces.
- Do not use on alcohol-sensitive materials (e.g. PLEXIGLAS®).
- PT 2 - Indoor:
 - Health care facilities and pharmaceutical and cosmetic industry, for example patient-near surrounding, working areas/desks, general equipment (with the exception of surfaces that come into contact with food).
- PT 4 - Health care facilities and in food industry, for example food preparation and handling in kitchens/restaurants.
- Authorisation number EU-0027671-000

Application notes:

- Surfaces should always be visibly clean before disinfection.
- Ready-to-use surface disinfectant at room temperature (20±2 °C).
- The entire surface to be disinfected is wetted by pouring or spraying from a short distance and subsequently thoroughly wiped with a cloth. The amount of product should be sufficient (max. 50 ml/m²) to keep the surface wet during the contact time.
- PT 2 - Application rate: Exposure time: - for the control of bacteria, yeasts and enveloped viruses: 60 sec - for the control of viruses (limited spectrum virucidal activity): 5 min.
 - Number and timing of application: A reasonable frequency of disinfection in a patient's room is 1-2 per day. Maximum number of applications is 6 per day. No safety intervals need to be considered between the application phases.
- PT 4 - Application rate: Minimum exposure time: for the control of bacteria and yeasts at 20°C: 60 sec.
 - Number and timing of application: the products can be used as often as necessary. A reasonable frequency in kitchens is 1-2 per day. No safety intervals need to be considered between the application phases.

Ingredients

Ingredients per 100g: 40g ISOPROPYL ALCOHOL, 25g PROPYL ALCOHOL

Sustainable development and production:



Product safety, storage and environmental instructions

Safety: For professional use only. For further information, see Safety Data Sheet. Use biocides safely. Always read the label and product information before use.

Storage: Store in dry, cool, well-ventilated place. Store in dry, cool, well-ventilated area. Keep container tightly closed. Keep out of direct sunlight. Recommended storage temperature: 0-30°C Do not store at temperatures below 0°C. Do not store near food, drink or animal feed. Keep away from combustible material.

Environmental: Disposal must be done according to official regulations. Do not empty into drains. Do not dispose of with domestic waste. Dispose of contents/container to an authorised waste collection point. Empty the packaging completely prior to disposal. When totally empty, containers are recyclable like any other packaging.

Application and dosing

Follow the instructions below.



concentrated
Disinfection.
 Suitable for areas with special hygiene requirements.



Surface disinfection:
 Wet the surface completely.



Toilet disinfection



concentrated
 Ideal for application on surfaces in kitchen areas.



Sales units:

- Order nr.: 716672 10 x 750 ml
- Order nr.: 714867 10 x 750 ml
- Order nr.: 714929 10 x 750 ml

Your trusted partner on site:



APESIN spray F



Alcohol-based rapid disinfection, liquid – perfume-free

Target organisms PT4	Dosage / Dilution	minimum exposure time
Bacteria	Ready-to-use	60 s
Yeasts		
PT 02 Target organism (s)	Dosage / Dilution	minimum exposure time
Bacteria	Ready-to-use	60 s
Yeasts		
Viruses (Virucidal activity against enveloped viruses)		
Viruses (Limited spectrum virucidal activity)	Ready-to-use	5 min
The results have been substantiated to the competent authority by testing the following standards: EN 1276, EN 1650, EN 13697, EN 13727, EN 13624, EN 13697 mod., EN 16615, EN 14476		
VAH -Certificate		minimum exposure time
Surface disinfection in medical areas, bactericidal and yeasticidal activity	with mechanical action dirty conditions	5 Min.

